

## APPENDIX "C"

1. Kosugi, S., et al (1993). Identification of thyroid-stimulating antibody-specific interaction sites in the N-terminal region of the thyrotropin receptor. *Molecular Endocrinology*, 7, 114-130.
- 5 2. Parma, J., et al (1993). Somatic mutations in the thyrotropin receptor gene cause hyperfunctioning thyroid adenomas. *Nature*, 365, 649-651.
3. Duprez, L., et al (1994). Germline mutations in the thyrotropin receptor gene cause non-autoimmune autosomal dominant hyperthyroidism. *Nature Genetics*, 7, 396-401.
- 10 4. Matus-Leibovitch, N., et al (1995). Truncation of the thyrotropin-releasing hormone receptor carboxy tail causes constitutive activity and leads to impaired responsiveness in *Xenopus* oocytes and AtT20 cells. *J. Biol. Chem.*, 270:3, 1041-1047.
5. Kosugi, S., et al (1995). Characterization of heterogenous mutations causing constitutive activation of the luteinizing hormone receptor in familial male precocious puberty. *Human Molecular Genetics*, 4 (No.2), 183-188.
- 15 6. Magnusson, Y., et al (1994). Autoimmunity in idiopathic dilated cardiomyopathy. *Circulation*, 89, 2760-2767.
7. Fu, M., et al (1994). Functional autoimmune epitope on alpha1-adrenergic receptors in patients with malignant hypertension. *Lancet*, 344, 1660-1663.
- 20 8. Samama, P., et al (1993a). A mutation-induced activation state of the B2-adrenergic receptor. *J. Biol. Chem.*, 268:7, 4625-36.
9. Kjelsberg, M.A., et al (1992). Constitutive activation of the alpha 1B-adrenergic receptor by all amino acid substitutions at a single site. *J. Biol. Chem.* 267, 1430-1433.
10. Ren, Q., et al (1993). Constitutively active mutants of the alpha2-adrenergic receptor. *J. Biol. Chem.*, 268, 16483-16487.
- 25 11. Burstein, E.S., et al (1996). Constitutive activation of chimeric m2/m5 muscarinic receptors and delineation of G-protein coupling selectivity domains. *Biochem Pharmacol*, 51:4, 539-44. Burstein, E.S., et al (1996). Amino acid side chains that define muscarinic receptor/G-protein coupling. Studies of the third intracellular loop. *J. Biol. Chem.*, 271:6, 2882-5.
- 30 12. Hasegawa, H., et al (1996). Two isoforms of the prostaglandin E receptor EP3 subtype different in agonist-independent constitutive activity. *J. Biol. Chem.*, 271:4, 1857-1860.
13. Nanevicz, T., et al (1996). Thrombin receptor activating mutations. *J. Biol Chem.*, 271, 702-706.
14. Boone, C., et al (1993). Mutations that alter the third cytoplasmic loop of the a-factor

receptor lead to a constitutive and hypersensitive phenotype. *Proc. Natl. Acad. Sci. (USA)*, 90:21, 9921-5.

15. Spiegel, A.M., et al (1995). Defects in G protein-coupled signal transduction in human disease. *Ann. Rev. Physiol.* 58, 143-170.
- 5 16. Seeman, P. (1993). Dopamine D4 receptors elevated in schizophrenia. *Nature*, 365, 441-445.
17. Mann, J., et al (1986). Increased serotonin<sub>2</sub> and beta-adrenergic receptor binding in the frontal cortices of suicide victims. *Arch. Gen. Psychiat.* 43, 954-959. Casey, C., et al (1996). Constitutively active mutant 5-HT<sub>2A</sub> serotonin receptors: inverse agonist activity of classical 5HT<sub>2A</sub> antagonists. *Soc. Neurosci. Abstracts* # 699.10
- 10 18. Barker, E.L., et al (1994). Constitutively active 5-hydroxytryptamine<sub>2C</sub> receptors reveal novel inverse agonist activity of receptor ligands. *J. Biol. Chem.*, 269:16, 11687-11690.
19. Maenhault, C., et al (1990). RCD8 codes for an adenosine A<sub>2</sub> receptor with physiological constitutive activity. *Biochem. Biophys. Res. Com.*, 173:3, 1169-1178.
- 15 20. Parfitt, A.M., et al (1996). Hypercalcemia due to constitutive activity of the parathyroid hormone (PTH)/PTH-related peptide receptor: comparison with primary hyperparathyroidism. *J. Clin. Endocr. Metab.*, 81, 3584-3588.
21. Lavlie, R., et al (1996). The Ca<sup>2+</sup>-sensing receptor gene (PCAR1) mutation T151M in isolated autosomal dominant hypoparathyroidism. *Hum Genet*, 98:2, 129-33.
- 20 22. Arvanitakis, L., et al (1997). Human herpesvirus KSHV encodes a constitutively active G-protein-coupled receptor linked to cell proliferation.
23. Liu, J., et al (1996). Molecular mechanisms involved in muscarinic acetylcholine receptor-mediated G protein activation studied by insertion mutagenesis. *J. Biol. Chem.*, 271:11, 6172-6178.
- 25 24. Prezeau, L., et al (1996). Changes in the carboxy-terminal domain of metabotropic glutamate receptor 1 by alternate splicing generate receptors with differing agonist-independent activity. *Mol. Pharmacol.*, 49, 422-429.
25. Alla, S.A., et al (1996). Extracellular domains of the bradykinin B<sub>2</sub> receptor involved in ligand binding and agonist sensing defined by anti-peptide antibodies. 271, 1748-1755.
- 30 26. Wang, Z., et al (1994). Constitutive  $\mu$  opioid receptor activation as a regulatory mechanism underlying narcotic tolerance and dependence. *Life Sciences*, 54:22, 339-350.
- 35 27. Tiberi, M. & Caron, M.G. (1994). High agonist-independent activity is a distinguishing feature of the dopamine D<sub>1B</sub> receptor subtype. *The J. Biol. Chem.* 269:45. 27925-

27931.

28. Robbins, L.S., et al (1993). Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function. *Cell*, 72, 827-834.
29. Eggerick, D., et al (1995). Molecular cloning of an orphan G-protein-coupled receptor that constitutively activates adenylate cyclase. *Biochem. J.* 389, 837-843.
30. Jakub/Eik, J., et al (1995). Constitutive activity of the M1-M4 subtypes of muscarinic receptors in transfected CHO cells and of muscarinic receptors in the heart cells revealed by negative antagonists. *FEBS Lett*, 377:2, 275-9.
31. Burstein, E.S., et al (1995). Constitutive activation of muscarinic receptors by the G-protein Gq. *FEBS Lett*, 363:3, 261-3.
32. Ter Lack, A., et al (1995). Modelling and mutation studies on the histamine H1-receptor agonist binding site reveal different binding modes for H1-agonism: Asp116 (TM3) has a constitutive role in receptor stimulation. *J. Computer-aided molecular design*, 9, 319-330.